

Role of fuziline in ameliorating cardiac damage

Cardiac damage and fuziline

Yasemin Hacanli, Ezhar Korkmaz Ersöz

Department of Cardiovascular Surgery, Faculty of Medicine, Harran University, Şanlıurfa, Turkey

Abstract

Modifiable and non-modifiable factors play a role in the occurrence of cardiovascular diseases, as well as free radicals. Free radicals are caused by a decrease in the antioxidant level or an increase in the amount of oxidant in the cells. In the studies, it has been suggested that the consumption of vegetables, fruits and foods with high antioxidant content is effective in preventing many diseases, especially phenolic compounds with antioxidant effect prevent diseases. Phenolic compounds are essential compounds for living things and form an important part of antioxidants. Many drugs and antioxidants are being tested for the treatment of cardiac damage. Fuzi (Radix Aconiti Lateralis Preparata), one of these antioxidant substances, is defined as a Chinese herb. It is a derived form of Aconitum carmichaelii Debx. Scientific studies on Fuzi; hypolipidemic activity, kidney protection, cardiogenic activity, immune system improvement, anti-arrhythmia, anti-aging and antineoplastic activity, etc. made for its features. Our aim in this review is to present the cardiovascular, neuropharmacological, antidepressant, anticancer, anti-inflammatory and analgesic effects of fuzi, which has antioxidant properties.

Keywords

Fuziline, Dobutamine, Antioxidant, Cardiovascular Diseases, Myocardial Damage

DOI:10.4328/ECAM.10041

Received : 2022-09-22

Accepted : 2022-10-11

Published Online : 2022-10-11

Printed Online : 2022-09-01

Eu Clin Anal Med 2022; 10(3):31-33

Corresponding Author: Yasemin Hacanli, Department of Cardiovascular Surgery, Faculty of Medicine, Harran University, Şanlıurfa, Turkey.

E-Mail: yaseminhacan@hotmail.com · **P:** +90 506 700 27 00 · **Corresponding Author ORCID ID:** <https://orcid.org/0000-0002-4427-8149>

How to cite this article: Yasemin Hacanli, Ezhar Korkmaz Ersöz. Role of fuziline in ameliorating cardiac damage. Eu Clin Anal Med 2022; 10(3):31-33

Introduction

Many factors play a role in the formation of cardiovascular diseases (CVD). These negative factors are; obesity, hypertension, diabetes etc. It consists of modifiable and non-modifiable factors such as genetic predisposition, age, and gender [1]. CVD that can result in death includes heart attack, heart failure, coronary heart disease, etc. It is possible to count heart diseases [2]. Also CVD, cancer etc. It is known that free radicals cause many diseases. Under optimal conditions, the harmony that exists between antioxidants and oxidants is disrupted by a decrease in the antioxidant level or an increase in the amount of oxidant [3]. The increase in free radical synthesis under deteriorated optimal conditions causes oxidative stress. It has been stated that oxidative stress is a serious factor in the pathophysiology of cardiac remodeling, which causes the formation and progression of heart failure (HF) [4].

Positive inotropic agent support is recommended to maintain end-organ activity and systemic perfusion in patients admitted to hospitals due to heart disease [5]. Agents that cause an increase in cardiac output by increasing the contractility of the heart muscle are called (positive) inotropic agents [6]. Dobutamine is widely used as a positive inotropic agent. Created from isoproterenol (ISO), it has less vascular and arrhythmogenic effects than other positive inotropic agents [7]. Today, various studies are carried out to investigate the effects in the diagnosis and treatment of CVD. In these studies, animal models that reflect the pathophysiology of CVD in humans are carried out. In CVD models created with dobutamine, it has been observed that oxidative stress generally increases and dobutamine creates immunosuppression on the immune system.

Studies have suggested that phenolic compounds with antioxidant effects prevent diseases [8]. Traditional Chinese medicine has historically been accepted as an alternative therapeutic method with less cost and less toxic effects against heart diseases and other diseases [9].

Phenolic compounds

Phenolic compounds are essential compounds for animals and humans and form an important part of antioxidants [10]. Phenolic compounds are named according to their constituent parts and ring forms [11]. In some studies, it has been reported that phenolic compound supplementation has efficacy in reducing or preventing the occurrence of oxidative stress-related diseases such as diabetes and CVD. It has also been found to play a role in preventing oxidative stress, increasing lipolysis, inhibiting adipocyte exchange and development, and reducing lipogenesis and inflammatory responses [12]. Phenolic compounds are divided into different groups as flavonoids, tocopherols and phenolic acids [13]. The largest group of phenolic compounds is flavonoids and ranks first in researches [14].

Flavonoids

Flavonoids have efficacy to reduce fibrosis or tissue damage and inhibit certain enzymes involved in inflammation. In vivo and in vitro studies have shown that flavonoids can stop the formation and progression of inflammatory disorders [15]. In addition, they have protective effects on CVD by suppressing lipoprotein oxidation [16]. Both phenolic and

flavonoid compounds have vasodilator, antithrombotic, antiallergic, anti-inflammatory, antimicrobial, antidiabetic properties. It is known that they exhibit antioxidant behaviors in preventing the formation of many diseases, especially CVD, cancer and Alzheimer's [17].

Fuziline

Molecular Formula: C₂₄H₃₉N₀₇

Molecular Weight: 453.27g/mol

Plant Origin: *Aconitum carmichaeli* debx [18].

Fuzi (*Radix Aconiti Lateralis Preparata*) [19], identified as a Chinese herb, is a derived form of *Aconitum carmichaelii* Debx [20]. Fuzi; Chinese wolfsbane, Chinese aconite, monkweed, Bushi in Japan and Kyeong-Po Buja in Korea etc. known by names [21]. Products derived from Fuzi are Yan - Fuzi, Heishunpian and Baifupian and their pharmacological effect has been accepted. There are 211 of the aconitum varieties in China. Among these, there are relatively 400 species in the northern temperate area [19]. Fuzi has 122 chemical components, mainly flavonoids, alkaloids, fatty acids and saponins [21]. Fuziline, one of the components of Fuzi; When compared with mesaconite, aconite and hypaconite diterpenoids in terms of negative effects, it has been seen that it has less negative effects and safer use [22]. To date, research on Fuzi modules has focused on the carbohydrate molecules of both aconitum alkaloids and this herbal medicine [23]. Recently, there has been research on the neuropharmacological activities of some of the substances in Fuzi [24]. Many researchers have investigated fuzi's pharmacological efficacy, detoxification methods and mechanisms, toxicity, etc. They have worked on the subject [25].

Discussion

Intensive scientific studies on fuzi; hypolipidemic activity, kidney protection, cardiotoxic activity, immune system improvement, anti-arrhythmia, anti-aging and antineoplastic activity, etc. was made for its properties [26]. In the literature review, many studies on the subject were found. Zhao et al. in their research on *aconitum carmichaelii*; Important analgesic effects of fuziline, hypaconitine, mesaconitine and neoline were determined. In the same study, apart from fuziline in mice; It has been determined that neoline, mesaconit and hypoconit have serious side effects. This situation revealed the reliability of fuziline. In addition, the maximum application dose of fuziline was stated as 1 g/kg . Considering this efficacy of Fuziline, the idea that it could be developed as an analgesic drug has emerged [27]. Liu et al. in a study on mice, they investigated the antidepressant properties of fuzi total alkaloid and showed that fuzi total alkaloid may have antidepressant properties in mice with ovaries removed [28]. Xu et al. in a study by Sprague-Dawley on rats, AWA [(Aconite water-soluble alkaloid extract and its content: fuziline, mesaconine, karakoline, hypaconine, neoline and songorine are active components of Fuzi and its side effects are low [29])] evaluated its effects on congestive heart failure due to upper abdominal aortic coarctation surgery. During this evaluation, hypertrophic effects including increased serum ANP, NT-pro BNP, TNF- α and IL-6 levels were detected in the model group, and it was reported that improvement was observed with AWA treatment [30]. Zhang et al. in their study on floppy-eared white rabbits in 2006, the effects on gastrointestinal microcirculation were investigated by preserving myocardial functions after myocardial ischemia reperfusion injury by using Shenfu injection consisting of shenfu and fuzi pharmaceutical compounds. At the end of the study, it was seen that Shenfu injection supports gastrointestinal microcirculation [31]. Fan et al. in their study, AST, lactate dehydrogenase (LDH), CK and CK-MB levels were examined in male Sprague Dawley rats in order to determine the myocardial damage caused by ISO in male Sprague Dawley rats, and these parameters were found to be higher in the ISO group and lower




Figure 1. Chemical Form of Fuziline [18].

in the fuziline and metoprolol groups. The protection of fuziline against myocardial damage has been clearly demonstrated and the idea that it can be used therapeutically has emerged [32].

Storch et al. demonstrated that fuzi alkaloids regulate cellular metabolism by acting on the electron transport system in mitochondria [33]. Lu et al. They suggested that Fuzi alkaloids could inhibit tumor formation by suppressing central carbon metabolism, HIF-1 and the PI3K/AKT/mTOR pathway [34]. Zhang et al. demonstrated that fuzi extract attenuated the amount of Treg, IL-10 and TGF- β synthesized in association with radiation, modulating immunity and inhibiting tumor development [35]. Li et al. reported in their study with network pharmacology analysis that anti-inflammatory activation by fuzi-containing Shenfu injection (SFI) could occur in a way associated with NF- κ B. Three compounds (including benzoyl-mesaconine, fuziline, and neoline) were the first SFI compounds reported to have NF- κ B inhibitory activity [36]. Sun et al. According to the pharmacokinetic studies on Sprague-Dawley rats by hydrophilic interaction liquid chromatography-electrospray ionization mass spectrometry method, fuziline was successfully administered and the clearance rate, half-life and bioavailability rate of fuziline were determined by oral administration of 4mg/kg in rats [22]. Gong et al. Pharmacokinetic study of fuziline by oral and intravenous administration in Beagle dog by UPLC-Q-TOF-MS method and it was reported that fuziline has acceptable accuracy and precision values [37].

Conclusion

According to the information obtained in researches on fuziline, its protective and supportive importance against CVDs, its anti-inflammatory, anticancer, analgesic and antidepressant properties come to the fore. Additional research on its biochemical activity and bioactive components is needed to better understand these effects. This review can be a resource for further basic research and clinical applications on fuziline.

Scientific Responsibility Statement

The authors of this article declare that they are responsible for the scientific content of the article, including the design of the study, data collection, analysis and interpretation, writing, preparation and scientific review of some or all content of the main line and approval of the final version.

Animal and human rights statement

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. No animal or human studies were carried out by the authors for this article.

Funding: None

Conflict of interest

None of the authors received any type of financial support that could be considered potential conflict of interest regarding the manuscript or its submission.

References

- Tanrıverdi B, Tetik ŞS. Aterosklerozun Patofizyolojisi ve Risk Faktörleri. *Marmara Pharmaceutical Journal*. 2017;21: 1-9.
- Sarrafzadegan N, Gotay C. CVD prevention in 2014: Advances in the prevention of cardiovascular disease. *Nat Rev Cardiol*. 2015;12:71-3.
- Karabulut H, Gülay MŞ. Serbest radikaller. *MAKÜ Sag. Bil. Enst. Derg*. 2016. 4(1):50-59.
- Tsutsui H, Kinugawa S, Matsushima S. Oxidative stress and heart failure. *Am J Physiol Heart Circ Physiol*. 2011;301(6):H2181-90.
- Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;62:147-239.
- Hensyl WR. *Stedman's medical Dictionary*. 25th ed. 1990, Baltimore: Williams and Wilkins.
- Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001; 8;345(19):1368-77.
- Priscilla HD, Prince PS. M. Cardioprotective effect of gallic acid on cardiac troponin-T, cardia marker enzymes, lipid peroxidation products and antioxidants in experimentally

- induced myocardial infarction in Wistar rats. *Chemico-Biological Interactions*. 2009;179: 118-124.
- Zhang K, Liu Y, Lin X, Yang J, Wu C. Assessment of Reproductive Toxicity and Genotoxicity of Aconiti Lateralis Radix Praeparata and its Processed Products in Male Mice. *J Ethnopharmacol*. 2021;275:114102.
- Gonçalves S, Moreira E, Grosso C, Andrade PB, Valentão P, Romano A. Phenolic profile, antioxidant activity and enzyme inhibitory activities of extracts from aromatic plants used in Mediterranean diet. *J Food Sci Technol*. 2017;54(1):219-227.
- Kolaç T, Gürbüz P, Yetiş G. Doğal Ürünlerin Fenolik İçeriği ve Antioksidan Özellikleri. *İnönü Üniversitesi Sağlık Hizmetleri Meslek Yüksekokulu Dergisi*. 2017;5(2).
- Rodríguez-Pérez C, Segura-Carretero A, Contreras DMM. Phenolic compounds as natural and multifunctional anti-obesity agents: A review. *Crit Rev Food Sci Nutr*. 2019;59(8):1212-1229.
- Hannan PA, Khan JA, Ullah I, Ullah S. Synergistic combinatorial antihyperlipidemic study of selected natural antioxidants; Modulatory effects on lipid profile and endogenous antioxidants. *Lipids Health Dis*. 2016;9:15(1):151.
- Baek SH, Cao L, Jeong JS, Kim HR, Nam J T, Lee GS. The Comparison of Total Phenolics, Total Antioxidant, and Anti-Tyrosinase Activities of Korean *Sargassum* Journal of food quality. 2021;1745-4557.
- Maleki SJ, Crespo JF, Cabanillas B. Anti-inflammatory effects of flavonoids. *Food Chem*. 2019;299:125124.
- Pietta PG. Flavonoids as antioxidants. *J Nat Prod*. 2000;63(7):1035-42.
- Comunian TA, Ravanfar R, De Castro IA, Dando R, Favaro-Trindade CS, Abbaspourrad A. Improving oxidative stability of echium oil emulsions fabricated by Microfluidics: effect of ionic gelation and phenolic compounds. *Food Chem*. 2017;233:125-134.
- Wu JJ, Guo ZZ, Zhu YF, Huang ZJ, Gong X, Li YH et al. A systematic review of pharmacokinetic studies on herbal drug Fuzi: Implications for Fuzi as personal-ized medicine. *Phytomed*. 2018;44:187-203.
- Xiao P, Wang F, Gao F, Yan L, Chen D, Liu Y. A pharmacophylogenetic study of Aconitum L. (Ranunculaceae) from China. *Acta Phytotax Sin*. 2006;01:1- 46.
- Chinese Pharmacopoeia Commission; Pharmacopoeia of the People's Republic of China. People's Medical Publishing House, Beijing, 2010;pp.177-178.
- Zhou G, Tang L, Zhou X, Wang T, Kou Z, Wang Z. A review on phytochemistry and pharmacological activities of the processed lateral root of Aconitum carmichaelii Debeaux. *J Ethnopharmacol*. 2015;3(160):173-93.
- Sun J, Zhang F, Peng Y, Liu J, Zhong Y, Wang G. Quantitative determination of diterpenoid alkaloid Fuziline by hydrophilic interaction liquid chromatography (HILIC)-electrospray ionization mass spectrometry and its application to pharmacokinetic study in rats. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2013;913-914:55-60.
- Wang BB, Wang JL, Yuan J, Quan Q-H, Ji R-F, Tan P, et al. Composition Analysis of Fuzi Polysaccharides by HPLC-MS(n) and Their Protective Effects on Schwann Cells Exposed to High Glucose. *Molecules*. 2016;21(11):1496.
- Farahani MS, Bahramsoltani R, Farzaei MH, Abdollahi M, Rahimi R. Plant-derived natural medicines for the management of depression: an overview of mechanisms of action. *Rev Neurosci*. 2015;26(3):305-21.
- Nyirimigabo E, Xu Y, Li Y, Wang Y, Agyemang K, Zhang Y. A review on phytochemistry, pharmacology and toxicology studies of Aconitum. *J Pharm Pharmacol*. 2015;67(1):1-19.
- Ren MY, Yu QT, Shi CY, Luo JB. Anticancer Activities of C18-, C19-, C20-, and Bis-Diterpenoid Alkaloids Derived from Genus Aconitum. *Molecules [Basel, Switzerland]*. *Molecules* 2017;22(2): 267.
- Zhao D, Shi Y, Zhu X, Liu L, Ji P, Long C, et al. Identification of Potential Biomarkers from Aconitum carmichaelii, a Traditional Chinese Medicine, Using a Metabolomic Approach. *Planta Med*. 2018;84(6-07):434-441.
- Liu L, Li B, Zhou Y, Wang L, Tang F, Shao D, et al. Antidepressant-like effect of Fuzi total alkaloid on ovariectomized mice. *J Pharmacol Sci*. 2012;120(4):280-7.
- Wang L, Ding JY, Liu XX, Tang MH, Chao RB, Wang FP. Identification of aminoalcohol-diterpenoid alkaloids in Aconiti Lateralis Radix praeparata and study of their cardiac effects. *Yao Xue Xue Bao*. 2014;49(12):1699-704.
- Xu X, Xie X, Zhang H, Wang P, Li G, Chen J, et al. Water-soluble alkaloids extracted from Aconiti Radix lateralis praeparata protect against chronic heart failure in rats via a calcium signaling pathway. *Biomed Pharmacother*. 2021;135:111184.
- Zhang XJ, Song L, Zhou ZG, Wang XM. Effect of shenfu injection on gastrointestinal microcirculation in rabbits after myocardial ischemia-reperfusion injury. *World J Gastroenterol*. 2006;12(27):4389-91.
- Fan CL, Yao ZH, Ye MN, Fu LL, Zhu GN, Dai Y, et al. Fuziline alleviates isoproterenol-induced myocardial injury by inhibiting ROS-triggered endoplasmic reticulum stress via PERK/eIF2 α /ATF4/Chop pathway. *J Cell Mol Med*. 2020;24(2):1332-1344.
- Storch A, Kaftan A, Burkhardt K, Schwarz J. 1-Methyl-6,7-dihydroxy-12,3,4-tetrahydroquinoline (Salsolinol) Is Toxic to Dopaminergic Neuroblastoma SH-Sy5y Cells via Impairment of Cellular Energy Metabolism. *Brain Res*. 2000;855(1):67-75.
- Lu CY, Ding ML, Cai SH, Xie H, Zhang W, Di LQ. Analysis of Alkaloids in Aconiti Lateralis Radix Praeparata and Prediction of Their Mechanism in the Treatment of Cancer. *J. Nanjing Univ. Traditional Chin. Med*. 2021;37(05):720-729.
- Zhang Q, Chen X, Luo Y, Ren H, Qiao T. Fuzi Enhances Anti-tumor Efficacy of Radiotherapy on Lung Cancer. *J Cancer*. 2017;8(19):3945-3951.
- Li P, Lv B, Jiang X, Wang T, Ma X, Chang N, Wang X, Gao X. Identification of NF- κ B inhibitors following Shenfu injection and bioactivity-integrated UPLC/Q-TOF-MS and screening for related anti-inflammatory targets in vitro and in silico. *J Ethnopharmacol*. 2016;194:658-667.
- Gong XH, Li Y, Li YX, Yuan A, Zhao MJ, Zhang RQ, et al. Development and Validation of an UPLC-Q-TOF-MS Method for Quantification of Fuziline in Beagle Dog After Intragastric and Intravenous Administration. *J Chromatogr Sci*. 2016;54(3):405-12.